



**IfBB**

Institute for Bioplastics  
and Biocomposites



# Biopolymers

facts and statistics

HOCHSCHULE  
HANNOVER  
UNIVERSITY OF  
APPLIED SCIENCES  
AND ARTS

–  
*Fakultät II  
Maschinenbau und  
Bioverfahrenstechnik*



1	INTRODUCTION AND BACKGROUND	4
2	PROCESS ROUTES	6
	GLOSSARY	7
2.1	Bio-based polyesters	8
2.1.1	Polylactic Acid (PLA)	8
2.1.2	Polyhydroxybutyrate (PHB)	10
2.1.3	Polybutylenesuccinate (PBS)	12
2.1.4	Polybutylenesuccinateadipate (PBSA)	15
2.1.5	Polytrimethyleneterephthalate (PTT)	18
2.1.6	Polyethyleneterephthalate (Bio-PET)	21
2.2	Bio-based polyolefins	24
2.2.1	Polyethylene (Bio-PE)	24
2.3	Bio-based polyamides (Bio-PA)	26
2.3.1	Homopolyamides	26
2.3.1.1	Bio-PA 6	26
2.3.1.2	Bio-PA 11	28
2.3.2	Copolyamides	29
2.3.2.1	Bio-PA 4.10 – Bio-PA 5.10 – Bio-PA 6.10	29
2.3.2.2	Bio-PA 10.10	30
2.4	Polyurethanes	32
2.5	Polysaccharide polymers	34
2.5.1	Cellulose-based polymers	34
2.5.1.1	Regenerated cellulose	34
2.5.1.2	Cellulose diacetate	35
2.5.2	Starch-based polymers	37
2.5.2.1	Thermoplastic starch (TPS)	37
2.5.2.2	Starch blends	38
3	MARKET DATA AND LAND USE FACTS	40
3.1	Global production capacities of bioplastics	42
3.2	Bioplastics production capacities by material type	43
3.3	Bioplastics production capacities by region	44
3.4	Bioplastics production capacities by market segment	45
3.5	Land use for bioplastics 2014 and 2019	46
	CONTACT	52

# 1

## INTRODUCTION AND BACKGROUND


The IfBB – Institute for Bioplastics and Biocomposites is a research institute within the Hochschule Hannover, University of Applied Sciences and Arts, which was established in 2011 to respond to the growing need for expert knowledge in the area of bioplastics. With its practice-oriented research and its collaboration with industrial partners, the IfBB is able to shore up the market for bioplastics and, in addition, foster unbiased public awareness and understanding of the topic.

As an independent research-led expert institution for bioplastics, the IfBB is willing to share its expertise, research findings and data with any interested party via the Internet, online and offline publications or at fairs and conferences. In carrying on these efforts, substantial information regarding market trends, processes and resource needs for bioplastics is being presented here in a concise format, in addition to the more detailed and comprehensive publication and “Engineering Biopolymers”<sup>1</sup>.

One of our main concerns is to furnish a more rational basis for discussing bioplastics and use fact-based arguments in the public discourse. Furthermore, “Biopolymers – facts and statistics” aims to provide specific, qualified answers easily and quickly for decision-makers in particular from public administration and the industrial sector. Therefore, this publication is made up like a set of rules and standards and largely foregoes textual detail. It offers extensive market-relevant and technical facts presented in graphs and charts, which means that the information is much easier to grasp. The reader can expect comparative market figures for various materials, regions, applications, process routes, agricultural land use or resource consumption, production capacities, geographic distribution, etc.

<sup>1</sup> Endres, H.-J.; Siebert-Raths, Andrea.  
Engineering Biopolymers:  
Markets, Manufacturing, Properties and  
Applications. Hanser 2011





In recent years, many new types of bioplastics have emerged and innovative polymer materials are pushing on the plastics market. All the same, bioplastics by no means constitute a completely new class of materials but rather one that has been rediscovered from among the large group of plastic materials.

The first polymer materials fashioned by human hands were all based on modified natural materials (e.g., casein, gelatine, shellac, celluloid, cellophane, linoleum, rubber, etc.). That means they were bio-based since petrochemical materials were not yet available at that time. Ever since the middle of the 20th century, these early bio-based plastics, with a few exceptions (cellulose and rubber-based materials), have almost been fully replaced by petrochemical materials.

By now, due to ecological concerns, limited petrochemical resources and sometimes new property profiles, bioplastics have undergone a remarkable revival and are taken more and more into focus by the general public, politics, the industrial sector and in particular the research community.

Of particular interest today are new types of bioplastics, which were developed in the past 30 years. The publication presented here refers to the so-called “New Economy” bioplastics as opposed to “Old Economy” bioplastics which indicate earlier materials developed before petrochemical bioplastics emerged, yet still exist on the market today (e.g., rubber, cellophane, viscose, celluloid, cellulose acetate, linoleum).

“New Economy” bioplastics divide up into two main groups. On the one hand, there are those biopolymers which have a new chemical structure virtually unknown in connection with plastics until a few years ago (e.g. new bio-based polyesters such as PLA), on the other hand so-called “drop-ins”, with the same chemical structure yet bio-based. The most prominent drop-ins at this point are bio-based PET (Bio-PET) and bio-based polyethylene (Bio-PE).

# 2

## PROCESS ROUTES

Process routes depict the manufacturing steps from the raw material to the finished product, specifying the individual process steps, intermediate products, and input-output streams. So they serve as a guide for all considerations and calculations around the production of bioplastics, in particular also with regard to their resource consumption.

The following methodical approach was chosen to establish the process routes:

The mass flows were first calculated using a molar method based on the chemical process, with the introduction of known rates and conversion factors. The routes so established were confirmed with polymer manufacturers and the industry. In so far as no loss rates due to the chemical processes or the process stages were included, the calculations were made basically assuming no losses. The mass flows show feedstock and resulting land requirements in ha for the production of one metric ton of bioplastics.

Feedstock requirements were calculated for the use of different crops. Yields of the most important crops and renewable raw materials used for feedstock are shown in the chart below. **Please note that the yields in this context refer to the crop itself, which contains the raw material for processing, and not to the harvested whole plant.**

FEEDSTOCK	CROP	RAW MATERIAL	AVERAGE CONTENT OF RAW MATERIAL	GLOBAL MEAN YIELD (Crop)	RESULTING AMOUNT (Raw material)	
CALCULATIONS			→	×	→	=
Corn	Maize kernel	Starch	70 %	6.5 t/ha	4.55 t starch/ha	
Potatoes	Potato tuber	Starch	18 %	21 t/ha	3.78 t starch/ha	
Wheat	Wheat grains	Starch	46 %	3.5 t/ha	1.60 t starch/ha	
Sugar cane	Sugar cane (without cane tops)	fermt. Sugar	13 %	70 t/ha	9.1 t sugar/ha	
Sugar beet	Beet (without leaves)	fermt. Sugar	16 %	52 t/ha	8.32 t sugar/ha	
Wood	Standing timber, residual wood	Cellulose	40 %	1.64 t atro/ha	0.66 t cellulose/ha	
Castor oil plant	Castor bean	Castor oil	40 %	1 t seeds/ha (given one harvest per year)	0.4 t oil/ha (given one harvest per year)	

## GLOSSARY

### ABBREVIATIONS USED:

bb = *bio-based*

fermt. = *fermentable*

SCA = *Succinic Acid*

BDO = *Butanediol*

PDO = *Propanediol*

PTA = *Purified Terephthalic Acid*

MEG = *Monoethylene Glycol*

PMDA = *Pentamethylene Diamine*

TMDA = *Tetramethylene Diamine*


HMDA = *Hexamethylene Diamine*

DMDA = *Decamethylene Diamine*

**red coloured** ressources have a petro-based origin


# 2.1 Bio-based polyesters

## 2.1.1 Polylactic Acid (PLA)



## PLA – Feedstock requirements (different feedstocks)

-  Sugar cane
-  Sugar beet
-  Corn
-  Wheat
-  Potato




## PLA – Land use in ha (different feedstocks)

-  Sugar cane
-  Sugar beet
-  Corn
-  Wheat
-  Potato





## 2.1.2 Polyhydroxybutyrat (PHB)




**\* Conversion Rates:**

Starch – Glucose 90 %


fermt. Sugar – PHB 35 %



## PHB – Feedstock requirements (different feedstocks)




## PHB – Land use in ha (different feedstocks)



## 2.1.3 Polybutylenesuccinate (PBS)

with bio-based Succinic Acid (PBS bb SCA)



**\* Conversion Rates:**

Starch – Glucose 90 %

fermt. Sugar – Succinic Acid 80 %




**IfBB**

Institute for Bioplastics  
and Biocomposites

## 2.1.3 Polybutylenesuccinate (PBS)

100% bio-based (PBS 100)



### \* Conversion Rates:


Starch – Glucose 90 %

fermt. Sugar – Succinic Acid 80 %




**IfBB**  
Institute for Bioplastics  
and Biocomposites

## PBS variations – Feedstock requirements (different feedstocks)




## PBS variations – Land use in ha (different feedstocks)



## 2.1.4 Polybutylenesuccinateadipate (PBSA)

with bio-based Succinic Acid (PBSA bb SCA)



### \* Conversion Rates:


Starch – Glucose 90 %

fermt. Sugar – Succinic Acid 80 %



## 2.1.4 Polybutylenesuccinateadipate (PBSA)

with bio-based Succinic Acid and Butanediol (PBSA bb SCA/BDO)




### \* Conversion Rates:

Starch – Glucose 90 %


fermt. Sugar – Succinic Acid 80 %



## PBSA variations – Feedstock requirements (different feedstocks)




## PBSA variations – Land use in ha (different feedstocks)





## 2.1.5 Polytrimethyleneterephthalate (PTT)

with bio-based Propanediol (PTT bb PDO)



**\* Conversion Rates:**

Starch – Glucose 90%

fermt. Sugar – Propanediol 40 %




**IfBB**


Institute for Bioplastics  
and Biocomposites

## 2.1.5 Polytrimethyleneterephthalate (PTT)


100% bio-based (PTT 100)



## PTT variations – Feedstock requirements (different feedstocks)




## PTT variations – Land use in ha (different feedstocks)




## 2.1.6 Polyethyleneterephthalate (Bio-PET)

with bio-based Ethanol (Bio-PET bb EtOH)




## 2.1.6 Polyethyleneterephtatlat (Bio-PET)


100% bio-based (Bio-PET 100)



## Bio-PET variations – Feedstock requirements (different feedstocks)




## Bio-PET variations – Land use in ha (different feedstocks)




# 2.2 Bio-based polyolefins


## 2.2.1 Polyethylene (Bio-PE)



## Bio-PE – Feedstock requirements (different feedstocks)



## Bio-PE – Land use in ha (different feedstocks)







# 2.3 Bio-based polyamides (Bio-PA)

## 2.3.1 Homopolyamides


### 2.3.1.1 Bio-PA 6



## Bio-PA 6 – Feedstock requirements (different feedstocks)




## Bio-PA 6 – Land use in ha (different feedstocks)



## 2.3.1 Homopolyamides

### 2.3.1.2 Bio-PA 11



<sup>1</sup> one harvest per year


**\* Conversion Rates:**

Ricinoleic Acid – Undecane Acid 50 %



## 2.3.2 Copolyamides

### 2.3.2.1 Bio-PA 4.10 – Bio-PA 5.10 – Bio-PA 6.10



<sup>1</sup> one harvest per year

• **Conversion Rates:**


Ricinoleic Acid – Sebacic Acid 60%



**IfBB**  
Institute for Bioplastics  
and Biocomposites

## 2.3.2 Copolyamides

### 2.3.2.2 Bio-PA 10.10




<sup>1</sup> one harvest per year

• **Conversion Rates:**


Ricinoleic Acid – Sebacic Acid 60 %





## Bio-PA – Feedstock requirements (feedstock castor oil)



## Bio-PA – Land use in ha (feedstock castor oil)




## 2.4 Polyurethanes




<sup>1</sup> one harvest per year

## Bio-PUR – Feedstock requirements (feedstock castor oil)



## Bio-PUR – Land use in ha (feedstock castor oil)






## 2.5 Polysaccharid polymers


### 2.5.1 Cellulose-based polymers (Cellulosics)

#### 2.5.1.1 Regenerated Cellulose




## 2.5.1 Cellulose-based polymers


### 2.5.1.2 Cellulose diacetate



Cellulosics – Feedstock requirements (feedstock wood)




Cellulosics – Land use in ha (feedstock wood)



## 2.5.2 Starch-based polymers


### 2.5.2.1 Thermoplastic starch (TPS)



\* Starch content 75 %

## 2.5.2 Starch-based polymers

### 2.5.2.2 Starch blends




\* Starch content 75 %


\*\* Ratio TPS/Polymer



## Starch-based polymers – Feedstock requirements (different feedstocks)



## Starch-based polymers – Land use in ha (different feedstocks)



# 3


## MARKET DATA AND LAND USE FACTS

As already mentioned in the introduction, the focus of attention is on “New Economy” bioplastics, including their position at the market. To give the reader an impression of the market share of these innovative and novel bioplastics:

when considering the most important Old Economy bioplastics with their global production capacity of 17 million tonnes annually, it turns out that the share of New Economy bioplastics is 10 times lower, i.e. 10 % of the market volume of all bio-based plastics (including the Old Economy bioplastics), with rising tendency.

By and large, Old and New Economy bioplastics (about 18.6 million tonnes) have a combined share of presently 6-7 % of the global plastics market. Given the anticipated market growth, especially of New Economy bioplastics, over a 5-year period, the market share of Old and New Economy bioplastics is expected to reach a maximum of 10 % of the global market for plastics within the next 5 years. The corresponding land use of Old and New Economy bioplastics is currently at approximately 15.6 million hectares, which is equivalent to only 0.3% of the global agricultural area or approximately 1% of the arable land. Comparing these figures reveals that New Economy bioplastics, which tend to be the only focus of interest in land use discussions, use up only 4 % of the area required for all bio-based plastics combined.

## Production capacities and land use Old and New Economy bioplastics



Even though global forecasts predict a rapidly growing market for these novel bioplastics in the next few years, the need for agricultural areas will be kept at a very low level. While the market for new bioplastics has been growing by around 15 % annually during the last three years and a sustained growth is anticipated in the future, it can be assumed that land use for New Economy bioplastics by 2019 (7.9 million tonnes), for example, will be as low as 0.03 % of the global agricultural area or about 0.1 % of the arable land. Regardless of the significant growth rates, it should be mentioned that the market share of these New Economy bioplastics is still hovering at less than 1 % of the global plastics market and is likely not to exceed 2 - 3 % in the near future. To make things even more compelling, it is a fact that bio-based plastics, even after multiple material usage, can still serve as an energy carrier. This means that additional crop lands, which are currently used for direct energy production, could be set aside for the production of bioplastics. Prior material usage of biomass, as in the case of bioplastics, still permits subsequent trouble-free energy recovery, whereas direct incineration of biomass (and also crude oil-based products!) precludes an immediate subsequent material usage. In this case, more arable land for plant cultivation is needed and consequently another photosynthesis process, in order to gain new resources once again as feedstock for material usage.


1 PLA, PHA, PTT, PBAT, Starch blends, Drop-Ins (Bio-PE, Bio-PET, Bio-PA) and other

2 material use excl. paperindustry

3 calculations include linseedoil only



# 3.1 Global production capacities of bioplastics



IfBB-Institute for Bioplastics and Biocomposites in cooperation with European Bioplastics e.V. / nova-Institute GmbH (Nov 2015)



**IfBB**  
Institute for Bioplastics  
and Biocomposites

## 3.2 Bioplastics production capacities by material type

Bioplastics production capacities 2014 (by material type)


**60.9%**  
bio-based/non-biodegradable



**39.1%**  
biodegradable



- 1 Biodegradable cellulose esters
- 2 Compostable hydrated cellulose foils
- 3 Bio-based content amounts 30%
- 4 Contains PBAT, PBS, PCL



IfBB-Institute for Bioplastics and Biocomposites in cooperation with European Bioplastics e.V. / nova-Institute GmbH (Nov 2015)

Bioplastics production capacities 2019 (by material type)


**83.6%**  
bio-based/non-biodegradable



**16.4%**  
biodegradable



- 1 Biodegradable cellulose esters
- 2 Compostable hydrated cellulose foils
- 3 Bio-based content amounts 30%
- 4 Contains PBAT, PBS, PCL




IfBB-Institute for Bioplastics and Biocomposites in cooperation with European Bioplastics e.V. / nova-Institute GmbH (Nov 2015)



**IfBB**  
Institute for Bioplastics  
and Biocomposites


# 3.3 Bioplastics production capacities by region

Global production capacities of bioplastics in 2014 (by region)



IfBB-Institute for Bioplastics and Biocomposites in cooperation with European Bioplastics e.V. / nova-Institute GmbH (Nov 2015)

Global production capacities of bioplastics in 2019 (by region)




IfBB-Institute for Bioplastics and Biocomposites in cooperation with European Bioplastics e.V. / nova-Institute GmbH (Nov 2015)



**IfBB**  
Institute for Bioplastics  
and Biocomposites


## 3.4 Bioplastics production capacities by market segment

Global production capacities of bioplastics 2014 (by market segment)



IfBB-Institute for Bioplastics and Biocomposites in cooperation with European Bioplastics e.V. / nova-Institute GmbH (Nov 2015)

Global production capacities of bioplastics 2019 (by market segment)



1 Contains regenerated cellulose and biodegradable cellulose ester 2 Bio-based content amounts to 30%

3 Contains durable starch blends, Bio-PC, Bio-TPE, Bio-PUR (except thermosets), Bio-PA, PTT


IfBB-Institute for Bioplastics and Biocomposites in cooperation with European Bioplastics e.V. / nova-Institute GmbH (Nov 2015)



# 3.5 Land use for bioplastics 2014 and 2019

For final land use calculation only the most commonly used crop was taken into consideration. Yield data from FAO statistics served as a basis for calculation (global, non-weighted average over the past 10 years). To calculate land use in this bottom-up approach, the producer-specific production capacities of a type of bioplastics were multiplied by the output data of the corresponding process routes.

In all of the calculations no allocation was made, which means land use was fully, by 100 %, allocated to the raw materials for bioplastics and not split up between various parallel side products such as proteins or straw in wheat. So this approach leads to a rather conservative estimate.



\* In relation to global agricultural area 2011

\*\* Also includes area growing permanent crops as well as approx. 1 % fallow land. Abandoned land resulting from shifting cultivation is not included.

Bioplastics	
<b>2014</b>	
683,000 ha	= 0.01 %*
<b>2019</b>	
1.4 million ha	= 0.03 %*

IfBB-Institute for Bioplastics and Biocomposites in cooperation with European Bioplastics e.V. / nova-Institute GmbH (Nov 2015)



**IfBB**  
Institute for Bioplastics  
and Biocomposites

IfBB – Institute for Bioplastics and Biocomposites

Hochschule Hannover | Heisterbergallee 12

D-30453 Hannover | Germany

Phone +49 511 9296-2268

Fax +49 511 9296-99 2268

E-Mail [info@ifbb-hannover.de](mailto:info@ifbb-hannover.de)

A large amount of additional information is also available on the  
IfBB website at [www.ifbb-hannover.de](http://www.ifbb-hannover.de).



**IfBB**  
Institute for Bioplastics  
and Biocomposites

